

Soluble endoglin: a biomarker or a protagonist in the pathogenesis of preeclampsia?

Lucia Perez-Roque, Jose M. Lopez-Novoa

Unidad de Fisiopatología Renal y Cardiovascular; Departamento de Fisiología y Farmacología, Universidad de Salamanca. 37007 Salamanca, Spain.

Instituto de Investigaciones Biomédicas de Salamanca, (IBSAL), 37007 Salamanca, Spain

Instituto "Reina Sofia" de Investigación Nefrológica, Fundación Renal Iñigo Alvarez de Toledo, 28034 Madrid, Spain

Received for publication: 28/07/2014

Accepted: 30/07/2014

INTRODUCTION

Pre-eclampsia is a pregnancy-specific disorder characterized by endothelial dysfunction, hypertension ($> 140/90$ mmHg) and proteinuria (> 300 mg/24hours) in the third trimester of pregnancy. It affects about 3-5% of all pregnancies worldwide^{1,2}. It remains a major cause of maternal mortality (15–46%) and is responsible for a 5-fold increase in perinatal mortality³. The World Health Organization has recognized the importance of pre-eclampsia by launching a global programme to fight pre-eclampsia-eclampsia (World Health Organization. Policy and Coordination Committee 16th meeting. Department of Reproductive Health and Research. Summary Medium-term Programme of Work 2004-2009 HRP/ PCC (16)/2003/8.1). In addition, women who have suffered pre eclampsia and their offspring are at greater risk of developing cardiovascular disease later in life⁴.

Several risk factors are associated for pre-eclampsia; these include nulliparity, multifetal gestations, previous history of pre-eclampsia, obesity, diabetes *mellitus*, vascular and connective tissue disorders like systemic lupus erythematosus and antiphospholipid antibodies. Severe pre-eclampsia can lead to the appearance of the HELLP (haemolysis, elevate liver function tests, low platelets) syndrome, the most critical variant of pre-eclampsia, and is

associated with a higher risk of maternal and neonatal adverse outcomes⁵.

At the present, there are no effective treatments for this disease, and delivery remains the only solution for maternal symptoms. In many cases, the newborn is often premature and there are significant perinatal morbidity and mortality among such babies.

Until recently, the molecular pathogenesis of pre-eclampsia was largely unknown, but recent observations support the hypothesis that altered expression of placental anti-angiogenesis factors are responsible for the clinical manifestation of this disease. Two anti-angiogenic factors soluble fms-like kinase 1 (sFlt1) and soluble endoglin (sEng) have been reported to have increased plasma concentration in the maternal circulation weeks before the onset of pre-eclampsia^{6,7} and decreased quickly after delivery⁸. The concentration of these factors in plasma is correlated with the severity of pre-eclampsia and bad prognosis. Thus, they have been proposed as useful biomarkers for the diagnostic and prognosis of pre-eclampsia^{9,10}.

Also, Gilbert *et al* showed that sEng and sFlt1 were increased in the serum of uteroplacental ischaemia animal model (RUPP)^{11,12}. sFlt1 is a soluble form of the VEGF receptor-1, a cell membrane receptor for VEGF (vascular endothelial growth factors) and PIGF

(placental growth factor-1), whereas sEng is the soluble form for a membrane co-receptor of several members of the TGF β superfamily. Both, sFlt1 and sEng, are secreted by the placenta^{13,14}.

■ PLACENTA IS THE ORIGIN OF PRE-ECLAMPSIA

Placenta plays a central role in pre-eclampsia as it is evidenced by rapid disappearance of disease symptoms after delivery and maintenance of these symptoms when placenta is retained. Abnormalities in the placental development have been studied in order to understand the early stage of this disease. The molecular basis for placental dysregulation of pathogenic factors it is not clear. The first event in the early gestation of pre-eclampsia is the abnormal placentation by alterations in the placenta vascular remodeling¹⁵. It has been reported unappropriated vascular remodeling of uterine spiral arteries by impaired invasion of these arteries by trophoblasts. On this way, the blood supply to the placenta is reduced to a level unable of providing adequate placental perfusion to sustain the growing fetus¹⁶ and generates a hypoxic environment. In humans, trophoblasts stem cells are differentiated into two cell types, villous trophoblasts and extravillous trophoblasts (EVTs). Earlier in pregnancy, invading trophoblasts, called interstitial EVT, migrate into the decidualized endometrium and endovascular EVT migrate along the lumen of spiral arterioles¹⁷. This EVT invasion creates the large-diameter and low-resistant vessels that carry blood to the placenta. After 12 weeks, endovascular EVT invade the uterine spiral arteries, replace the endothelial cells, and participate in the degradation of tunica media smooth muscle cells. This remodeling of the spiral arteries is essential for proper placental perfusion to sustain fetal growth. Pre-eclampsia is considered to be caused by poor placentation that can be explained by the shallow invasion by trophoblast until 20 weeks of gestation. This impairment is closely related with a failure of vascular remodeling. The consequence is shallow spiral arteries that contribute to the hypoxic environment in the placenta^{18,19}. Therefore, the hypoxic environment has been postulated as a critical signal that initiates the pathogenic process of pre-eclampsia²⁰. There are a lot of evidence supporting the central role of hypoxia in pre-eclampsia. It was reported that the placentas of pre-eclamptic women and models of placental hypoxia show similar gene expression²¹. In addition, reductions in utero placental blood flow in animal models induced

hypertension, proteinuria and endothelial dysfunction, which are the typical sign of this disease²². This relationship between uteroplacental ischaemia and pre-eclampsia was demonstrated in several animals, such as dogs²³, rabbits^{24,25}, primates^{26,27} and, more recently, it has been established a reduced uterine perfusion pressure model in the pregnant rat (RUPP) in which the utero placental perfusion is reduced by approximately 40% and shows many features of pre-eclampsia²⁸⁻³².

The possibility that sFlt1 and sEng are a key pathogenic link between placental pathology and maternal endothelial damage provides hope that these substances, in addition to serving as biomarkers of pre-eclampsia, could also play some role in the pathogenesis of the disease and, thus, could possibly also be effective therapeutic targets for the disease. In this editorial we will focus on sEng, in order to analyze if it is only a biomarker or if it could also be a protagonist in the pathogenesis of pre-eclampsia.

■ ENDOGLIN

Endoglin (Eng), also called CD105, is a glycosylated transmembrane protein that acts as an auxiliary receptor of several members of the TGF- β superfamily of proteins¹³. Endoglin is highly expressed in endothelial cells and syncytiotrophoblast^{33,34}. Although the most studied physiological effect of Eng is regulating angiogenesis [1], we have reported that Eng plays a major role in the regulation of vascular tone¹³. It has been described that the vasodilator effect of Eng is mediated by regulating endothelial nitric oxide synthase (eNOS) expression and NO synthesis³⁵. It has been proposed that Eng plays a major role in pre-eclampsia³⁶. In this review, we will describe the mechanisms of Eng overexpression and sEng release in pre-eclampsia, as well as the possible role of sEng in hypertension, proteinuria and abnormal placentation, all of them characteristic of pre-eclampsia.

■ Mechanisms of placental endoglin release in pre-eclampsia

Hypoxia seems to be involved in increased endoglin expression in the placenta

Endoglin is overexpressed in placenta during pre-eclampsia. Eng mRNA expression in chorionic villi

at 11-week's gestation were significantly higher in women that will develop pre-eclampsia. These findings suggest that this increase could be implicated in the early pre-eclampsia³⁷. The development of the placenta during the early normal pregnancy is carried out in an environment of low oxygen tension until 10 or 12 weeks of gestation. In that moment, the intervillous space is filled to maternal blood and the oxygen levels begin to increase. High level of endoglin is found in the early stage of gestation when the oxygen is low, but in the later stage the levels of endoglin is decreased^{38,39}. However, in pre-eclamptic placentas the amount of endoglin is always elevated. In trophoblasts obtained from placentas of pre-eclampsia patients, the release of sEng in hypoxia conditions (2% O₂ tension) was also higher than in trophoblast from normal pregnant women⁴⁰. These facts confirm that the hypoxic environment induces endoglin expression and plays an important role in the pathogenesis of pre-eclampsia⁴¹⁻⁴³. The mechanism explaining high endoglin expression in the presence of placental hypoxia is based in the fact that endoglin gene promoter has a hypoxia responsive element (HRE) region. HIF-1, in cooperation with Sp1 and the coactivator CBP/P300 is bound to this region stimulating endoglin expression. This stimulatory effect is further enhanced when the HIF-1 α /Sp1 complex collaborates with the TGF β /Smad3 signaling pathway⁴¹.

On the other hand, there are data suggesting that the increase in Eng expression observed in pre-eclampsia can also be mediated, at least in part, by HIF-1 α , probably through activation of the TGF- β 3 pathway^{43,44}. Tal R. *et al* have demonstrated the effects of HIF-1 overexpression in pregnant mice using adenovirus which express a constitutively active and stabilized HIF-1 α . They observed an elevation of sFlt1 and sEng in the plasma of these mice⁴⁵.

Increased oxidative stress can be involved in increased Eng expression by the placenta

A crosstalk between hypoxia and oxidative stress has been described by several authors. Under hypoxic conditions, reactive oxygen species (ROS) are produced by a flavoprotein-containing NAD(P)H or by mitochondria as a mechanism that contributes to stabilize HIF-1 α and consequently mediates the transcriptional response to hypoxia⁴⁶. Reactive oxygen

species generated at mitochondrial complex III stabilize hypoxia-inducible factor-1 α during hypoxia, nuclear translocation and transcriptional activation by a mechanism of O₂ sensing⁴⁶. Thus, hypoxic conditions may favor the generation of ROS in pre-eclampsia. Indeed, oxidative stress has been shown to play a pivotal role in the etiopathogenesis of pre-eclampsia²⁰. Furthermore, it has been demonstrated that a decrease in endogenous antioxidant factors, such as the antioxidant hemoxygenase-1 (HO-1), superoxide dismutase and glutathione peroxidase, may contribute to the increase release of sEng⁴⁷⁻⁴⁹.

Several studies have demonstrated high levels of LDL in the serum from women with pre-eclampsia and these LDL show a higher susceptibility to oxidation giving oxidized LDL (oxLDL) rich in oxysterols⁵⁰.

Endogenously oxygenated cholesterol derivatives, such as 24(S)-hydroxycholesterol or 22(R)-hydroxycholesterol (oxysterols), are detected at relatively high levels in the placenta⁵¹. Oxysterols, a product of cholesterol oxidation, can play a role mediating sEng release in pre-eclampsia. Oxysterols have been identified as endogenous agonists of liver X receptors (LXRs). LXRs are ligand-activated transcription factors, which have two isoform: LXR α and LXR β . LXR α is present in tissues with high lipid metabolism, such as liver, intestine, adipose tissue and kidney, whereas LXR β is ubiquitously expressed⁵². It has been suggested a role of LXRs in the reproductive system and a potential application of LXR agonists in the treatment of reproductive pathologies⁵³. It has been reported the association of the LXR β activation and pre-eclampsia. Mouzart *et al.* showed that the single nucleotide polymorphisms rs2695121 in LXR β gene was strongly associated with pre-eclampsia. In contrast, they did not find a significant association between the polymorphisms in LXR α gene and this disease⁵⁴. On the other hand, Henry-Berger *et al.* have shown that endoglin is one of the target gene of LXR. Endoglin has six putative response elements (LXRE) in its proximal promoter region. The heterodimer LXR/RXR is able to bind them and stimulate its transcription. It has been reported that endoglin was up-regulated by administration of LXR agonists in human choriocarcinoma JAR cells⁵⁵. Therefore, the LXR activation is able to up-regulate the inappropriate expression and function of endoglin and contribute with the complication of pregnancy⁵⁶.

Mechanisms involved in the release of sEng in the placenta

In pre-eclamptic women, the high levels of sEng seems be derived from membrane isoform of the placental cells¹⁴. The release of sEng occurs by proteolytic cleavage of the membrane isoform. It is known that this proteolytic cleavage is mediated by membrane metalloproteinases, which are increased in pregnancy. Hawinkels *et al*, reported that metalloproteinase-14 (MMP₁₄ or MT₁-MMP) is involved in the release of sEng⁵⁷. Kaitu'u-Lino *et al*. treated the syncytiotrophoblast cell line (BeWo) with metalloproteinase inhibitor, GM6001, and observed a significant decreased in sEng release. The GM6001 inhibits other MMPs besides MMP₁₄, so they administered MMP-13 or MMP-2/9 specific inhibitors, and did not observe differences in sEng release compared with its controls⁵⁸. Furthermore Kaitu'u-Lino, *et al.*, also demonstrated that there are other metalloproteinase present in placenta, such as MMP-16,-24,-25, but they were not associated with pre-eclampsia. Furthermore, MMP-17 and -15 were also up-regulated in pre-eclampsia, but neither of them were involved in the increased release of sEng^{59,60}.

Recently, a study from our group demonstrated that hypoxia induces sEng shedding in cultured human trophoblasts (JAR cells) and this release was increased by ROS. This study also demonstrated that oxysterols are involved in the sEng shedding through LXR pathway and that oxysterols and activators of LXR pathway increase cell surface expression and activity of MMP₁₄ and inhibit the TIM-3 expression which is a MMP₁₄ inhibitor⁵⁵. Thus, we can suggest that oxysterols induce sEng release by increasing cell surface endoglin⁵⁶, and by increasing MMP₁₄ and inhibit its inhibitor TIMP-3⁵⁵. Experiments *in vivo* have also demonstrated that administration of LXR agonist or oxysterols induced the increase in plasma levels of sEng in mice supporting the hypothesis that the LXR activation increases sEng release⁵⁵.

On the other hand, Lamarca *et al*, in the utero-placental ischaemia animal model (RUPP), have reported high levels of antibodies to angiotensin receptor type I, AT₁-AA. They also demonstrated that the increase of AT₁-AA induces the rise of TNF- α , which stimulates the release of sEng and sFlt₁⁶¹.

Thus, from our own results and those of the literature we can hypothesize that soluble endoglin

release is based on placental hypoxia, which causes oxidative stress and the formation of oxysterols. As a consequence, there is an increase in placental levels of membrane endoglin and the metalloprotease MMP-14, which cleaves membrane endoglin releasing the soluble form, thus increase plasma levels of this form (Fig. 1).

■ Possible effects of endoglin in pre-eclampsia


Effects of sEng in arterial pressure

The possible role of sEng in pre-eclampsia is not elucidated yet. It has been hypothesized that sEng plays its antiangiogenic and pro-hypertensive effects in pre-eclampsia due to its ability to bind circulating TGF β . TGF β , in the presence of membrane Eng, produce vasodilatation mediated by eNOS overexpression⁶². Thus, the binding of TGF- β by sEng could prevent the interaction of these TGF β molecules with the cell-surface Eng and, consequently, its normal effects in the endothelium⁶³. By this mechanism, sEng leads abolishes TGF β -dependent eNOS activation, thus leading to vasoconstriction¹⁴. Several studies reported that decreased NO availability is one of the principal causes of endothelial dysfunction in pre-eclampsia and is involved in haemodynamics changes characteristics of pre-eclampsia^{64,65}. However, this hypothesis needs to be adequately confirmed. Other authors have suggested that sEng alone cannot scavenge TGF β for which it has very low affinity, and suggest that sEng binds with high affinity to BMP₉, which stimulates secretion from endothelial cells of the vasoconstrictor endothelin-1, also implicated in endothelial cell stabilization⁶⁶. It remains to be determined if scavenging of circulating BMP₉ by sEng is important in pre-eclampsia and regulation of hypertension. It should be noted that administration of LXR agonist or oxysterols, in addition to inducing the increase in plasma levels of sEng, also induces hypertension in mice, whereas when sEng release was prevented by a blocking peptide, hypertension was not observed⁵⁵. Furthermore, male mice transgenic for sEng and with high plasma sEng levels also show hypertension⁵⁵. In the RUPP model in rats, the hypertension is associated to high plasma levels of sEng¹¹. Thus, all these data suggest that soluble endoglin plays a major role in the pathogenesis of pre-eclampsia-associated hypertension.

Figure 1

Mechanisms proposed for soluble endoglin (sEng) release and effects of sEng in the pathogenesis of pre-eclampsia.

sEng release is based on placental hypoxia (with increase of cytosolic levels of HIF-1), which causes oxidative stress and the formation of oxysterols. This situation activates the LX receptors and NF κ B pathway. As a consequence, there is an increase in placental levels of membrane endoglin and the metalloprotease MMP-14, which cleaves membrane endoglin releasing the soluble form, thus increasing plasma levels of this endoglin form. High plasma levels of endoglin decrease vessel relaxation ability, thus contributing to placental hypoperfusion (and, consequently, to placental hypoxia) and increased peripheral resistance (and, consequently, increased arterial pressure). sEng also impairs normal trophoblast invasion and normal placentation. In addition, sEng could also alter renal glomerular filtration membrane, thus inducing proteinuria.



Effects of sEng in proteinuria

Classically, glomerular endothelial injury was considered responsible for the renal impairment present in pre-eclampsia. Proteinuria, another major characteristic of pre-eclampsia, has been explained by “endotheliosis” of glomerular capillaries, an alteration characterized by endothelial cell swelling and loss of endothelial fenestrae in these capillaries. It has been demonstrated that neutralization of TGF- β by neutralizing TGF-beta1 antibodies results in loss of glomerular fenestrae and proteinuria, thus suggesting that the glomerular endotheliosis lesions of pre-eclampsia are at least partially due to the placenta-derived sEng, which block the TGF-beta/endoglin signaling, respectively, causing the loss of glomerular endothelial cell fenestrae, cell swelling and proteinuria⁶⁷.

Podocytes are modified glomerular epithelial cells that cover the outside surface of glomerular capillaries, and play a major role in regulating glomerular permeability to proteins. Recent evidence suggests that appearance of podocytes in urine is associated with the appearance and prognosis of pre-eclampsia⁶⁸. There are some data suggesting that podocytes are crucial in explaining the loss of filtration capacity of the pre-eclamptic kidney⁶⁹. Also, recent studies from our laboratory have demonstrated that sEng is able to prevent leukocyte binding to endothelium by interfering in the binding of leukocyte integrins to the RGD domain of endothelial endoglin⁷⁰. Thus, it can be hypothesized that by a similar mechanism, sEng is able to reduce the binding of the podocytes to glomerular basal membrane (GBM) by impairing the binding of integrin $\alpha 3 \beta 1$ in the podocyte to

the GBM component laminin-521, one of the major mechanism for podocyte attachment⁷¹. This could induce the detachment of podocytes from the GBM, as these cells are subjected to a great force by the fluid that is filtered through the wall of the glomerular capillaries.

Effects of sEng in placentation

As above described, endoglin is greatly expressed in syncytiotrophoblast and extravillous trophoblast (EVT). It was shown that the inappropriate levels of endoglin could induce a wrong trophoblast differentiation through a negative regulation by TGF β 1 and TGF β 3⁷². Studies have demonstrated that overexpression of endoglin decreased cell migration and invasion in mouse fibroblast or endothelial cells. Endoglin also has an important role in trophoblast biology⁴³, through its ligands, TGF β 1 and TGF β 3⁷². TGF β signaling is involved in several inhibitory signals, such as antiproliferative, antimigratory and anti-invasive signals. Furthermore, it is demonstrated that LXR β activation reduced EVT invasion and migration through TGF β pathway⁷³.

In addition, sEng can be also involved in inadequate placentation. It has been reported that sEng reduces the invasive ability of cytotrophoblasts through regulation of the expression of MMP-2 and MMP-9, metalloproteases necessary for a correct trophoblast invasion⁷⁴. It has been reported that the invasion and vascular remodeling under hypoxia were significantly reduced in autophagy-deficient EVT cells compared with wild-type EVT cells. Interestingly, sEng suppressed EVT invasion by inhibiting autophagy. This effect of sEng was recovered by TGF- β 1 treatment in a dose-dependent manner. A high dose of sEng inhibited the vascular construction by EVT cells and endothelial cells, whereas a low dose of sEng inhibited the replacement of endothelial cells by EVT cells⁷⁵. Although the evidences are very limited, we can hypothesize that sEng is involved in the inadequate placentation, which is the primary cause of pre-eclampsia. Thus, we can hypothesize, in the basis of the above described, that increased plasma levels of sEng are not only an early biomarker of pre-eclampsia but are also involved in the pathogenesis of the disease (defective placentation, hypertension and proteinuria) (Fig. 1)

■ CONCLUSION AND PERSPECTIVES

In summary, placenta plays the central role in pre-eclampsia. The first event in early gestation is an abnormal placentation by alterations in vascular remodeling. The triggering of the hypoxia pathway induces an oxidative stress in the placenta. The oxidative stress and high levels of cholesterol in pre-eclamptic women induce an increased production of oxysterols in maternal plasma. This leads to activation of the oxysterols receptors LXR, which induces sEng release, and increased plasma levels of sEng contributes to the clinical manifestation of pre-eclampsia: hypertension and proteinuria, and contributes to defective placentation, thus creating a vicious circle that amplifies the disease. A schematic representation of these mechanisms is shown in Fig. 1. According with these results, it can be speculated that one of the therapies for pre-eclampsia should be the reduction of sEng shedding. We have hypothesized that inhibition of the enzymatic activity of MMP₁₄ could prevent sEng release. We have described a peptide (P583), which corresponds to the aminoacids 583-590 of endoglin sequence. This sequence includes the MMP₁₄ cleavage site GL and interferes with its proteolytic activity. It was proven *in vitro* that P583 inhibits the sEng shedding. Administration of p583 to mice treated with LXR agonist or oxysterols reduced plasma levels of sEng and reverted the appearance of hypertension in these mice⁵⁵. These studies suggest that a therapeutic strategy that inhibits sEng release or eliminates sEng from the circulation could be effective in preventing pre-eclampsia, a disease without treatment at present.

Acknowledgments

The authors thank Dr. Carmelo Bernabeu, (*Centro de Investigaciones Biológicas*, CSIC, Madrid, Spain) for his advice with the endoglin projects. Studies from the authors' laboratories have been supported by grants from *Ministerio de Economía y Competitividad of Spain* (SAF2010-15881 and SAF2013-45784-R), *Junta de Castilla y Leon* (Grant GR100), *Red de Investigación Cooperativa en Enfermedades Renales* (Red-inRen, RDo6/0016/0013 and RD12/0021/0032) and *Instituto "Reina Sofía" de Investigación Nefrológica, Fundación Renal Iñigo Álvarez de Toledo*. Red-inRen is an initiative of the *Instituto de*

Salud Carlos III (ISCIII) of Spain supported by FEDER funds. LP-R is supported by a fellowship from *Junta de Castilla y Leon* (ORDEN EDU/346/2013).

Conflict of interest statement: None declared

References

- Bernabeu C, Lopez-Novoa JM, Quintanilla M. The emerging role of TGF-beta superfamily coreceptors in cancer. *Biochim Biophys Acta* 2009;1792(10):954-973.
- Eiland E, Nzerue C, Faulkner M. Preeclampsia 2012. *J Pregnancy* 2012; doi: 10.1155/2012/586578.
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol* 2009;33(3):130-137.
- Smith GC, Pell JP, Walsh D. Pregnancy complications and maternal risk of ischaemic heart disease: a retrospective cohort study of 129,290 births. *Lancet* 2001;357(9273):2002-2006.
- Rachdi R, Fekih MA, Massoudi L, et al. [HELLP syndrome. Epidemiological, nosological and prognostic aspects]. *Rev Fr Gynecol Obstet* 1993;88(4):230-235.
- Levine RJ, Lam C, Qian C, Yu KF, et al. Soluble endoglin and other circulating antiangiogenic factors in preeclampsia. *N Engl J Med* 2006;355(10):992-1005.
- Troisi R, Braekke K, Harsem NK, Hyer M, Hoover RN, Staff AC. Blood pressure augmentation and maternal circulating concentrations of angiogenic factors at delivery in preeclamptic and uncomplicated pregnancies. *Am J Obstet Gynecol* 2008;199(6):653 e1-10.
- Purwosunu Y, Sekizawa A, Farina A, et al. Evaluation of physiological alterations of the placenta through analysis of cell-free messenger ribonucleic acid concentrations of angiogenic factors. *Am J Obstet Gynecol* 2008;198(1):124 e1-7.
- Lopez-Novoa JM. Soluble endoglin is an accurate predictor and a pathogenic molecule in pre-eclampsia. *Nephrol Dial Transplant* 2007;22(3):712-714.
- Crovetto F, Figueras F, Triunfo S, et al. Added value of angiogenic factors for the prediction of early and late preeclampsia in the first trimester of pregnancy. *Fetal Diagn Ther* 2014;35(4):258-266.
- Gilbert JS, Gilbert SA, Arany M, Granger JP. Hypertension produced by placental ischemia in pregnant rats is associated with increased soluble endoglin expression. *Hypertension* 2009;53(2):399-403.
- Gilbert JS, Babcock SA, Granger JP. Hypertension produced by reduced uterine perfusion in pregnant rats is associated with increased soluble fms-like tyrosine kinase-1 expression. *Hypertension* 2007;50(6):1142-1147.
- Lopez-Novoa JM, Bernabeu C. The physiological role of endoglin in the cardiovascular system. *Am J Physiol Heart Circ Physiol* 2010;299(4):H959-974.
- Venkatesha S, Toporsian M, Lam C, et al. Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 2006;12(6):642-649.
- Pijnenborg R, Dixon G, Robertson WB, Brosens I. Trophoblastic invasion of human decidua from 8 to 18 weeks of pregnancy. *Placenta* 1980;1(1):3-19.
- Powe CE, Levine RJ, Karumanchi SA. Preeclampsia, a disease of the maternal endothelium: the role of antiangiogenic factors and implications for later cardiovascular disease. *Circulation* 2011;123(24):2856-2869.
- Zhou Y, Fisher SJ, Janatpour M, et al. Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for successful endovascular invasion? *J Clin Invest* 1997;99(9):2139-2151.
- Lyall F. Priming and remodelling of human placental bed spiral arteries during pregnancy—a review. *Placenta* 2005;26 Suppl A:S31-36.
- Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental development by oxygen tension. *Science* 1997;277(5332):1669-1672.
- Al-Gubory KH, Fowler PA, Garrel C. The roles of cellular reactive oxygen species, oxidative stress and antioxidants in pregnancy outcomes. *Int J Biochem Cell Biol* 2010;42(10):1634-1650.
- Soleymanlou N, Jurisica I, Nevo O, et al. Molecular evidence of placental hypoxia in preeclampsia. *J Clin Endocrinol Metab* 2005;90(7):4299-4308.
- Granger JP, Alexander BT, Llinas MT, Bennett WA, Khalil RA. Pathophysiology of preeclampsia: linking placental ischemia/hypoxia with microvascular dysfunction. *Microcirculation* 2002;9(3):147-160.
- Abitbol MM, Pirani CL, Ober WB, Driscoll SG, Cohen MW. Production of experimental toxemia in the pregnant dog. *Obstet Gynecol* 1976;48(5):537-548.
- Abitbol MM, Gallo GR, Pirani CL, Ober WB. Production of experimental toxemia in the pregnant rabbit. *Am J Obstet Gynecol* 1976;124(5):460-470.
- Losonczy G, Brown G, Venuto RC. Increased peripheral resistance during reduced uterine perfusion pressure hypertension in pregnant rabbits. *Am J Med Sci* 1992;303(4):233-240.
- Combs CA, Katz MA, Kitzmiller JL, Brescia RJ. Experimental preeclampsia produced by chronic constriction of the lower aorta: validation with longitudinal blood pressure measurements in conscious rhesus monkeys. *Am J Obstet Gynecol* 1993;169(1):215-223.
- Hodari AA. Chronic uterine ischemia and reversible experimental "toxemia of pregnancy". *Am J Obstet Gynecol* 1967;97(5):597-607.
- Alexander BT, Kassab SE, Miller MT, et al. Reduced uterine perfusion pressure during pregnancy in the rat is associated with increases in arterial pressure and changes in renal nitric oxide. *Hypertension* 2001;37(4):1191-1195.
- Alexander BT, Rinewalt AN, Cockrell KL, Massey MB, Bennett WA, Granger JP. Endothelin type a receptor blockade attenuates the hypertension in response to chronic reductions in uterine perfusion pressure. *Hypertension* 2001;37(2 Pt 2):485-489.
- Alexander BT, Cockrell K, Cline FD, Llinas MT, Sedeek M, Granger JP. Effect of angiotensin II synthesis blockade on the hypertensive response to chronic reductions in uterine perfusion pressure in pregnant rats. *Hypertension* 2001;38(3 Pt 2):742-745.
- Crews JK, Herrington JN, Granger JP, Khalil RA. Decreased endothelium-dependent vascular relaxation during reduction of uterine perfusion pressure in pregnant rat. *Hypertension* 2000;35(1 Pt 2):367-372.
- Granger JP, LaMarca BB, Cockrell K, et al. Reduced uterine perfusion pressure (RUPP) model for studying cardiovascular-renal dysfunction in response to placental ischemia. *Methods Mol Med* 2006;122:383-392.
- Gougos A, St Jacques S, Greaves A, et al. Identification of distinct epitopes of endoglin, an RGD-containing glycoprotein of endothelial cells, leukemic cells, and syncytiotrophoblasts. *Int Immunol* 1992;4(1):83-92.
- Cheifetz S, Bellon T, Cales C, et al. Endoglin is a component of the transforming growth factor-beta receptor system in human endothelial cells. *J Biol Chem* 1992;267(27):19027-19030.
- Toporsian M, Gros R, Kabir MG, et al. A role for endoglin in coupling eNOS activity and regulating vascular tone revealed in hereditary hemorrhagic telangiectasia. *Circ Res* 2005;96(6):684-692.
- Oujo B, Perez-Barriocanal F, Bernabeu C, Lopez-Novoa JM. Membrane and soluble forms of endoglin in preeclampsia. *Curr Mol Med* 2013;13(8):1345-1357.
- Farina A, Sekizawa A, De Sanctis P, et al. Gene expression in chorionic villous samples at 11 weeks' gestation from women destined to develop preeclampsia. *Prenat Diagn* 2008;28(10):956-961.
- Rodesch F, Simon P, Donner C, Jauniaux E. Oxygen measurements in endometrial and trophoblastic tissues during early pregnancy. *Obstet Gynecol* 1992;80(2):283-285.
- Burton GJ, Caniggia I. Hypoxia: implications for implantation to delivery—a workshop report. *Placenta* 2001;22 Suppl A:S63-65.
- Gu Y, Lewis DF, Wang Y. Placental productions and expressions of soluble endoglin, soluble fms-like tyrosine kinase receptor-1, and placental growth factor in normal and preeclamptic pregnancies. *J Clin Endocrinol Metab* 2008;93(1):260-266.
- Sanchez-Elsner T, Botella LM, Velasco B, Langa C, Bernabeu C. Endoglin expression is regulated by transcriptional cooperation between the hypoxia and transforming growth factor-beta pathways. *J Biol Chem* 2002;277(46):43799-43808.
- Caniggia I, Winter JL, Adriana and Luisa Castellucci Award lecture 2001. Hypoxia inducible factor-1: oxygen regulation of trophoblast differentiation in normal and preeclamptic pregnancies—a review. *Placenta* 2002;23 Suppl A:S47-57.
- Yinon Y, Nevo O, Xu J, et al. Severe intrauterine growth restriction pregnancies have increased placental endoglin levels: hypoxic regulation via transforming growth factor-beta 3. *Am J Pathol* 2008;172(1):77-85.
- Caniggia I, Mostachfi H, Winter J, et al. Hypoxia-inducible factor-1 mediates the biological effects of oxygen on human trophoblast differentiation through TGFbeta(3). *J Clin Invest* 2000;105(5):577-587.
- Tal R, Shaish A, Barshack I, et al. Effects of hypoxia-inducible factor-1alpha overexpression in pregnant mice: possible implications for preeclampsia and intrauterine growth restriction. *Am J Pathol* 2010;177(6):2950-2962.
- Chandel NS, McClintock DS, Feliciano CE, et al. Reactive oxygen species generated at mitochondrial complex III stabilize hypoxia-inducible factor-1alpha during hypoxia: a mechanism of O2 sensing. *J Biol Chem* 2000;275(33):25130-25138.
- Tam Tam KB, Lamarca B, Arany M, et al. Role of reactive oxygen species during hypertension in response to chronic antiangiogenic factor (sFlt-1) excess in pregnant rats. *Am J Hypertens* 2011;24(1):110-113.
- Wang Y, Walsh SW. Antioxidant activities and mRNA expression of superoxide dismutase, catalase, and glutathione peroxidase in normal and preeclamptic placentas. *J Soc Gynecol Investig* 1996;3(4):179-184.
- Cudmore M, Ahmad S, Al-Ani B, et al. Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. *Circulation* 2007;115(13):1789-1797.
- Wakatsuki A, Ikenoue N, Okatani Y, Shinohara K, Fukaya T. Lipoprotein particles in preeclampsia: susceptibility to oxidative modification. *Obstet Gynecol* 2000;96(1):55-59.
- Qiu C, Phung TT, Vadachkoria S, Muiy-Rivera M, Sanchez SE, Williams MA. Oxidized low-density lipoprotein (Oxidized LDL) and the risk of preeclampsia. *Physiol Res* 2006;55(5):491-500.
- Repa JJ, Mangelsdorf DJ. The role of orphan nuclear receptors in the regulation of cholesterol homeostasis. *Annu Rev Cell Dev Biol* 2000;16:459-481.
- Bourdeau A, Dumont DJ, Letarte M. A murine model of hereditary hemorrhagic telangiectasia. *J Clin Invest* 1999;104(10):1343-1351.
- Mouzat K, Mercier E, Polge A, et al. A common polymorphism in NR1H2 (LXRbeta) is associated with preeclampsia. *BMC Med Genet* 2011;12:145.

55. Valbuena-Diez AC, Blanco FJ, Oujó B, *et al.* Oxysterol-induced soluble endoglin release and its involvement in hypertension. *Circulation* 2012;126(22):2612-2624.
56. Henry-Berger J, Mouzat K, Baron S, *et al.* Endoglin (CD105) expression is regulated by the liver X receptor alpha (NR1H3) in human trophoblast cell line JAR. *Biol Reprod* 2008;78(6):968-975.
57. Hawinkels LJ, Kuijper P, Wiercinska E, Verspaget HW, Liu Z, Pardali E, *et al.* Matrix metalloproteinase-14 (MT1-MMP)-mediated endoglin shedding inhibits tumor angiogenesis. *Cancer Res* 2010;70(10):4141-4150.
58. Kaitu'u-Lino TJ, Palmer KR, Whitehead CL, Williams E, Lappas M, Tong S. MMP-14 is expressed in preeclamptic placentas and mediates release of soluble endoglin. *Am J Pathol* 2012;180(3):888-894.
59. Kaitu'u-Lino TJ, Palmer K, Tuohy L, Ye L, Tong S. MMP-15 is upregulated in preeclampsia, but does not cleave endoglin to produce soluble endoglin. *PLoS one* 2012;7(6):e39864.
60. Kaitu'u-Lino TJ, Tuohy L, Ye L, Palmer K, Skubisz M, Tong S. MT-MMPs in pre-eclamptic placenta: relationship to soluble endoglin production. *Placenta* 2013;34(2):168-173.
61. Zhou CC, Irani RA, Zhang Y, *et al.* Angiotensin receptor agonistic autoantibody-mediated tumor necrosis factor-alpha induction contributes to increased soluble endoglin production in preeclampsia. *Circulation* 2010;121(3):436-444.
62. Santibanez JF, Letamendia A, Perez-Barriocanal F, *et al.* Endoglin increases eNOS expression by modulating Smad2 protein levels and Smad2-dependent TGF-beta signaling. *J Cell Physiol* 2007;210(2):456-468.
63. Luft FC. Soluble endoglin (sEng) joins the soluble fms-like tyrosine kinase (sFlt) receptor as a pre-eclampsia molecule. *Nephrol Dial Transplant* 2006;21(11):3052-3054.
64. Lowe DT. Nitric oxide dysfunction in the pathophysiology of preeclampsia. *Nitric Oxide* 2000;4(4):441-458.
65. Shesely EG, Maeda N, Kim HS, *et al.* Elevated blood pressures in mice lacking endothelial nitric oxide synthase. *Proc Natl Acad Sci USA* 1996;93(23):13176-13181.
66. Gregory AL, Xu G, Sotov V, Letarte M. Review: the enigmatic role of endoglin in the placenta. *Placenta* 2014;35 Suppl:S93-99.
67. Ballemann BJ. Contribution of the endothelium to the glomerular permselectivity barrier in health and disease. *Nephron Physiol* 2007;106(2):p19-25.
68. Craici IM, Wagner SJ, Weissgerber TL, Grande JP, Garovic VD. Advances in the pathophysiology of pre-eclampsia and related podocyte injury. *Kidney Int* 2014;86(2):275-285.
69. Henao DE, Saleem MA. Proteinuria in preeclampsia from a podocyte injury perspective. *Curr Hypertens Rep* 2013;15(6):600-605.
70. Rossi E, Sanz-Rodríguez F, Eleno N, *et al.* Endothelial endoglin is involved in inflammation: role in leukocyte adhesion and transmigration. *Blood* 2013;121(2):403-415.
71. Sachs N, Sonnenberg A. Cell-matrix adhesion of podocytes in physiology and disease. *Nat Rev Nephrol* 2013;9(4):200-210.
72. Caniggia I, Taylor CV, Ritchie JW, Lye SJ, Letarte M. Endoglin regulates trophoblast differentiation along the invasive pathway in human placental villous explants. *Endocrinology* 1997;138(11):4977-4988.
73. Mo J, Fang SJ, Chen W, Blobel GC. Regulation of ALK-1 signaling by the nuclear receptor LXRbeta. *J Biol Chem* 2002;277(52):50788-50794.
74. Dong W, Xu QX, Han YH, Yin LR. [Inhibitory effects of soluble endoglin on invasive ability of cultured cytotrophoblasts of first trimester of pregnancy]. *Sheng Li Xue Bao* 2011;63(3):267-271.
75. Nakashima A, Yamanaka-Tatematsu M, Fujita N, *et al.* Impaired autophagy by soluble endoglin, under physiological hypoxia in early pregnant period, is involved in poor placentation in preeclampsia. *Autophagy* 2013;9(3):303-316.

Correspondence to:

José M. López-Novoa, Ph.D.

Departamento de Fisiología y Farmacología; Edificio Departamental; Campus Miguel de Unamuno; 37007 Salamanca (Spain)

E-mail: jmlnovoa@usal.es